



# Perinatal bacterial infection: screening of vertical transmitted infections

Maria Teresa Neto MD, PhD

Neonatal Intensive Care Unit  
Hospital de Dona Estefânia, Centro Hospitalar de Lisboa Central, EPE,  
Faculdade de Ciências Médicas, Universidade Nova de Lisboa  
Portugal



UNIVERSIDADE NOVA DE LISBOA







**Hospital Dona Estefânia, Lisbon. Founded in 1860**

# Summary

- The rational for screening
- Why some countries have a non-screening policy
- Risk factors: a very useful complement of screening
- Conclusions



# The burden of the problem

- Incidence of early-onset neonatal infection may vary between 0.98 to 1.2 or 1.3/1000 live births (*Stoll B, Kuhn P, Lin C-Y*)
- Lethality vary between 1.8% and 16% ((Kuhn, Stoll).
- Mortality depends on gestational age and isolates: 30% in newborns 25-28 weeks; 33% in *E. Coli* vs 9% for GBS (*Barbara Stoll, 2011*)

# Early-onset bacterial neonatal infection

Caused by bacteria colonising the birth canal

- Gram positive isolates:  
*Group B Streptococcus*  
but also *Streptococcus pneumoniae*,  
*Enterococcus*, *Listeria*
- Gram negative isolates:  
*Enterobacteriaceae* -  
*E. coli* but also  
*Proteus* spp, *Klebsiella*  
spp, *Haemophilus*

- Vergnano S et al. *Arch Dis Child Fetal Neonatal* Ed 2011 UK
- Kuhn P et al. *Paediatric and Perinatal Epidemiol* 2010 France
- Stoll BJ et al. *Pediatrics* 2011 - USA
- Lin CY et al. *Pediatr Neonatol* 2011 - Taiwan.
- Al-Ta'ar A et al. *International J Infectious Dis* 2011 Kuwait

# Early-onset bacterial neonatal infection

- In the past Enterobacteriaceae, mostly *E. coli*, were accepted as the most common cause of early-onset (EOS) neonatal bacterial infection
- In the 1970s GBS infections emerged as the leading cause of EOS and meningitis (and *E. Coli* was forgotten?)




# Portuguese National Prevalence Study

## One day surveillance - 2010

	Ner of patients	Bacteria
Newborns in the postnatal ward	567	<i>Enterococcus faecalis</i> – 1
Newborns in NICU	287	<i>E. coli</i> – 6 GBS – 2 <i>Streptococcus pneumoniae</i> – 1 <i>Proteus mirabilis</i> – 1
Patients >28 days Infections in patients admitted from home	20270	<i>E.coli</i> - 320 GBS – 22 <i>Listeria</i> -2

# Incidence of early-onset infection caused by GBS and *E. coli* /1000 births





# Sepsis caused by possible mother-acquired bacteria in newborn infants admitted to NICUs

Portuguese surveillance system data

	2008	2009	2010	Total
Patients	5369	5058	4911	15338
GBS	34 6.3/1000	27 5.3/1000	20 4.1/1000	81 (5/1000)
<i>E. coli</i>	37 6.9/1000	39 7.7/1000	24 4.9/1000	100 (7/1000)
<i>Listeria</i>	6	0	4	10 (0.7/1000)

# GBD early-onset infection

- Since 1992 AAP guidelines advise screening and prevention of GBS vertical transmission aiming to decrease early-onset infection.
- These guidelines were adopted by several countries and denied by others
- No studies on screening and prophylaxis for *E. coli* are known

# The Portuguese studies

## PPSU

- Group B streptococcal disease in Portuguese infants younger than 90 days

Enrolling only septic newborn infants with positive cultures in sterile fluids \* (2001-2004)

- Group B streptococcal disease - the hidden cases

Enrolling newborn infants with proven and possible GBS infection (2006-2007)

\* *Neto MT Arch. Dis. Child. Fetal Neonatal Ed. 2008;93;90-93*

\*\* *Neto MT Non-published data*



## Compared data


	First study	Second study
Incidence of early-onset proven infection	0.44/1000LB	0.22/1000LB
Early-onset infection	83%	66%

# Incidence


Per complete years, in the two studies

Proven infection only

Casualty or effects of screening and prophylaxis?



# GBS and *E. coli* positive blood cultures in Portuguese NICU in two different periods (in % of admitted newborn infants)





# Conclusions of two important studies on universal GBS screening and prophylaxis

- Rates of GBS infection declined but reflect a continued burden of disease. GBS continues the most frequent pathogen in term infants and *E. Coli* the most frequent in preterm infants

- *Stoll BJ et al. Pediatrics 2011 - USA.*

- GBS screening and prophylaxis is effective in decreasing the incidence of GBS EOS; however an increase in EOS caused by *E. Coli* was noted.

*Lin CY et al. Pediatr Neonatol 2011 - Taiwan.*

Is there a reason to screen? Yes!

However, even today, against all grades of evidence, some countries do not have national guidelines

Why?

# Bias related to carriers

- Colonization is intermittent
- Prevalence of carriers varies with geographic areas making obsolete national guidelines. In Portugal it is about 30% in the North and 12% in the South\*

*\*Neto MT, 2009*



# Bias related to screening

- Screening should be done by 35/36 gestational week - preterm delivery is excluded
- Vaginal and rectal swabs should be done - many women only have a vaginal swab - false negative
- Specification on request should be done - false negative
- Transport and culture should be appropriate - false negative
- Once a national guideline is implemented it should be accomplished all over the country with uniform high quality

# Bias related to prophylaxis

- Time before birth is needed
- Penicillin is the antibiotic of choice.
- GBS resistance to some of the alternatives have been reported - 10% to erythromycin in a Portuguese study\*
- Early-onset infection may occur in newborn infants whose mothers were given correct prophylaxis

*\* Exposto F. Non-published data*

# Newborn infants with early-onset GBS infection

## Screening and prophylaxis

	<b>Neto MT</b>	<b>Stoll B</b>
Screened mothers	61%	58%
Negative screening	28%	81% in term and 26% in PT
Intrapartum Antibiotics	28%	53% all EOS



# Newborns with early-onset infection and mother's prophylaxis

Proven and possible infection - n=57

Positive	Complete prophylaxis	1 dose of antibiotics	None
19	7* (37%)	9	3

\* One infant with positive blood culture

Bias related to those conditions  
supposed to protect that do not

Proven and possible GBS early-onset infection	Caesarean section
57	24 (42%)*

\* 14 (58%) with positive blood culture

# Screening and prophylaxis

## Summarizing confounding factors

### Reasons for non-screening policies

- Newborn infants with early-onset infection born to supposed negative mothers
- Newborn infants with early-onset infection born to mothers with complete prophylaxis
- Newborn infants with early-onset infection born by caesarean section



If screening has so many problems  
should prophylaxis be based on risk  
factors?

# Newborn infants with early-onset proven infection Known risk factors

Non-screened mothers with risk factors	57 (29%)
Term newborn infants with early-onset infection and risk factors	35/160 (22%) *

Conclusion: Prophylaxis based on risk factors would have missed 71% of all newborn infants and 78% of term newborns with early-onset infection

P.J.Steer and J Plumb found that only 60% of newborn infants with EOGBS disease had risk factors apparent at labour

*(Semin Fetal Neonatal Med 2011;16: 254-8)*



# Common-sense proposals

- Screening should be correctly performed - rectal and introit swabs, selective medium
- Women should know the meaning of carrier state in order to go to the maternity on time to start prophylaxis
- Antibiotics should be started as soon as possible instead of awaiting for a schedule

# Common-sense proposals

- GBS carrier state should not be considered the only main risk for early-onset neonatal infection.
- Other risk factors have to be considered even if there is a negative GBS screening - maternal fever, prolonged rupture of membranes, laboratory signs of mother's infection deserve attention and prophylaxis or treatment
- In any case we know some cases will be missed

# Final conclusions

- Screening and intrapartum antibiotics, result in a significant decrease of EOGBS disease. However GBS infection still exists. Better screening and prophylaxis and avoidance of missed opportunities to prevent neonatal infection are desirable
- *E. coli* is the second most common bacteria causing EOS and its importance should not be disregarded mainly in preterm infants
- Monitoring of pathogens causing EOS continues an important issue
- Adding screening to risk factors seems to be a valuable policy to improve prevention of EOS